



Topazio

Unique disease resistance package



Topazio has a unique disease resistance package now being offered to the industry! The first hybrid grey zucchini that has resistance to squash leaf curl virus, in addition to zucchini yellow mosaic virus and powdery mildew. Topazio's fruit are slightly lighter in color than Amatista. The plants are very upright, allowing for easier harvesting than older varieties. Fruit has a classical zucchini grey fruit color and shape. Topazio is a definite benefit for the times of the year when squash leaf curl, zucchini yellow mosaic virus and powdery mildew are the limiting factors to zucchini production.

T R I A L D A T A	
Approx. Days to Maturity	45
Average Length (in.)	4 - 5
Average Diameter (in.)	1.75
Shape	Tear - drop
Fruit Color	Whitish green with speckles
Plant Type	Upright, vigorous plant.
Disease Resistance	IR: ZYMV; SCLV; Sf

* See Back Side for Disease Resistance Descriptions




Topazio



Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. ROGERS® is a registered trademark of a Syngenta Group Company. Syngenta Seeds, Inc., P.O. Box 4188, Boise, ID 83711-4188, U.S.A. www.rogersadvantage.com



KEY TO RESISTANCE ABBREVIATIONS FOR SQUASH

CMV	Cucumber mosaic caused by <i>Cucumber mosaic virus</i>
Sf	Powdery mildew caused by <i>Sphaerotheca fuliginea</i>
WMV	Watermelon mosaic caused by <i>Watermelon mosaic virus</i>
ZYMV	Zucchini yellows caused by <i>Zucchini yellow mosaic virus</i>
(HR)	High Resistance: describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
(IR)	Intermediate Resistance: describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
	The VIP seal denotes Value-added, Innovation and Performance

Pathogen races are indicated to the right of the colon (example: Xcv: 1, 2, 3 = Bacterial spot caused by races 1, 2 and 3 of *Xanthomonas campestris* pv. *vesicatoria*). In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.

Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield, quality, and level of claimed pest and pathogen resistances, are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield, quality, and level of claimed pest and pathogen resistances. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. Making Superior Vegetables a Reality™ is a trademark of Syngenta Group Company. ROGERS® is a registered trademark of Syngenta Group Company.